# Introducing





Auvi-Q is the **first-and-only** compact epinephrine auto-injector with **audio and visual cues** that guide patients and caregivers step-by-step through the injection process during a life-threatening allergic reaction. Auvi-Q is **easy-to-use**, is the **size and shape of a credit card and the thickness of a smartphone**, and fits comfortably in a pocket or small purse.

Auvi-Q is indicated for the emergency treatment of life-threatening allergic reactions, or anaphylaxis, in people who are at risk for or who have a history of these reactions.

Auvi-Q contains **epinephrine**, a well-established, first-line treatment for severe, life-threatening allergic reactions. Anaphylaxis, a serious allergic reaction that happens quickly and may cause death, can occur as a result of exposure to allergens including nuts, shellfish, dairy, eggs, insect bites, latex and medication, among other allergens.

#### Auvi-Q Features

Auvi-Q's breakthrough design features include:

- •) Audio and visual cues that guide users through the injection process. This includes voice-guided instructions with alert lights that signal when the injection is complete.
- •) Unique compact size and shape. Auvi-Q is 3½ inches tall, 2 inches wide and a half inch thick.
- •) Press-and-hold action with 5-second hold time
- •) Retractable Needle Mechanism designed to help prevent accidental needle sticks post-injection. Users will not see the needle before, during or after administration.

Auvi-Q should only be injected into your outer thigh. Do not inject into buttock or intravenously.

#### How To Get Auvi-Q

Auvi-Q is available by prescription in two dosage strengths:

- •) Auvi-Q 0.3mg is intended for patients who weigh 66 pounds or more
- •) Auvi-Q 0.15mg is intended for patients who weigh 33 66 pounds
- •) Studies in pediatric patients weighing less than 15kg (33lbs) have not been conducted

Each prescription contains two **0.3mg** or **0.15mg** single-dose devices and a training device. The training device is designed to familiarize patients and caregivers with how to use Auvi-Q in an allergic emergency.

It is recommended that patients at risk for anaphylaxis carry two doses of epinephrine with them all of the time. More than two sequential doses of epinephrine should only be administered under direct medical supervision.

Go to www.Auvi-Q.com for full prescribing information, and to see and hear Auvi-Q.

### Indication

Auvi-Q (epinephrine injection, USP) is used to treat life-threatening allergic reactions (anaphylaxis) in people who are at risk for or have a history of these reactions.

## Important Safety Information

Auvi-Q is for immediate self (or caregiver) administration and does not take the place of emergency medical care. Seek immediate medical treatment after use. Please see additional Important Safety Information on page 2.

#### Additional Important Safety Information

Auvi-Q<sup>TM</sup> should only be injected into your outer thigh. DO NOT INJECT INTO BUTTOCK OR INTRAVENOUSLY. If you accidently inject Auvi-Q into any other part of your body, seek immediate medical treatment. Epinephrine should be used with caution if you have heart disease or are taking certain medicines that can cause heart-related (cardiac) symptoms.

If you take certain medicines, you may develop serious life-threatening side effects from epinephrine. Be sure to tell your doctor all the medicines you take, especially medicines for asthma. Side effects may be increased in patients with certain medical conditions, or who take certain medicines. These include asthma, allergies, depression, thyroid disease, Parkinson's disease, diabetes, high blood pressure, and heart disease.

The most common side effects may include increase in heart rate, stronger or irregular heartbeat, sweating, nausea and vomiting, difficulty breathing, paleness, dizziness, weakness or shakiness, headache, apprehension, nervousness, or anxiety. These side effects go away quickly, especially if you rest.

Talk to your healthcare professional to see if Auvi-Q is right for you.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.



