

# Iron: Is Too Much Harmful?

## IMPACT OF EXCESS IRON IN THE BODY

### IRON IN THE BODY

Iron is an essential element in the human body mostly found in red blood cells. Iron helps cells "breathe" by carrying oxygen to cells and tissues, and is essential to giving the body energy and having a properly functioning immune system<sup>1</sup>.

Most people get iron from the food they eat<sup>2</sup>.



ANIMAL SOURCES



FORTIFIED PRODUCTS




OTHER SOURCES

Iron circulates through the body continuously and any unused iron is stored for future use<sup>3</sup>.

- People with too little iron in their body can develop iron deficiency, which causes anemia<sup>1</sup>.
- People who accumulate too much iron in their body may develop chronic iron overload<sup>3</sup>.

When the body's iron capacity is exceeded and the body cannot get rid of it, iron builds up—first in the liver, and eventually in the heart. This condition is called chronic iron overload<sup>3</sup>.



## HOW IRON ENTERS THE BODY

People can develop chronic iron overload through:

- **Blood transfusions** required for managing many chronic health conditions such as **sickle cell disease, thalassemia, and myelodysplastic syndromes (MDS)**<sup>3,4</sup>.
- **Increased absorption** through the stomach and intestines. This can happen even in patients who do not receive regular blood transfusions, such as non-transfusion-dependent thalassemia (NTDT) patients<sup>3</sup>. Increased iron absorption in NTDT patients is triggered by the body's need for more red blood cells.


### FOOD



10-15% ABSORBED  
1-2 MG TAKEN IN

A healthy person absorbs and releases about 1-2 mg of iron each day, or about 10-15% of the iron in a normal diet<sup>1,2</sup>.


### INTESTINAL ABSORPTION



20% ABSORBED  
3-4 MG TAKEN IN

A person with non-transfusion-dependent thalassemia (NTDT) absorbs twice as much iron from their food as a normal person, amounting to 3-4 mg of iron each day, or about 20% of the iron in a normal diet<sup>5,1</sup>.

### BLOOD TRANSFUSION



BLOOD TRANSFUSION  
200 MG TAKEN IN

A person who receives blood transfusions absorbs an average of 200 mg of extra iron from each unit of blood transfused<sup>6</sup>. As few as 20 units of blood (10 in children) can lead to chronic iron overload<sup>3</sup>.

#### REFERENCES

1. Muñoz M, Villar I, García-Erce JA. An update on iron physiology. *World J Gastroenterol.* 2009 Oct 7;15(37):4617-26.
2. National Institutes of Health. Dietary Supplement Factsheet: Iron. Accessed June 25, 2013 from <http://ods.od.nih.gov/factsheets/Iron-HealthProfessional/>.
3. Shander A, Cappellini MD, et al., Iron overload and toxicity: the hidden risk of multiple blood transfusions. *Vox Sanguinis* 2009; 97, 185-197.
4. Kohgo Y, Ikuta K, Ohtake T, Torimoto Y, Kato J. Body iron metabolism and pathophysiology of iron overload. *Int J Hematol.* 2008 Jul;88(1):7-15.
5. Musallam K. Iron overload in non-transfusion-dependent thalassemia. *Thalassemia Reports* 2013;3:34-36.
6. Thalassemia International Federation. Guidelines for the clinical management of thalassemia. Nov. 2008.
7. Yen A. W. et al., Revisiting Hereditary Hemochromatosis: Current Concepts and Progress. *The American Journal of Medicine* 2006; 119, 391-399.
8. Taher AT. Age-related complications in treatment-naive patients with thalassaemia intermedia. *B J Haematol.* 2010;150:480-97.