



OVERVIEW

Alimera Sciences, Inc. (NASDAQ: ALIM), based in Alpharetta, GA, is a pharmaceutical company that specializes in the research, development and commercialization of prescription ophthalmic pharmaceuticals. Alimera Sciences Limited, a subsidiary of Alimera, is based in the United Kingdom and is the headquarters of Alimera's European operations. The company also operates subsidiaries in Germany and Portugal. Presently, the company is focused on diseases affecting the back of the eye, or retina.

ILUVIEN® (fluocinolone acetonide intravitreal implant) 0.19 mg is the company's lead product. ILUVIEN is a multiyear intravitreal (in the vitreous of the back of the eye) nonbioerodable implant, which is a sustained drug delivery system that provides a continuous, daily submicrogram level of fluocinolone acetonide lasting for 36 months.

ILUVIEN is approved in the United States to treat diabetic macular edema in patients who have been previously treated with a course of corticosteroids and did not have a clinically significant rise in intraocular pressure. ILUVIEN is also approved in Austria, Belgium, Denmark, Finland, France, Germany, Ireland, Italy, Luxembourg, Netherlands, Norway, Portugal, Spain, Sweden and the United Kingdom where it is indicated for the treatment of vision impairment associated with chronic diabetic macular edema, considered insufficiently responsive to available therapies. ILUVIEN is in the national licensing phase in two additional European countries, the Czech Republic and Poland. It is available commercially in the U.S., Germany, Portugal and the United Kingdom.

MANAGEMENT TEAM

- Dan Myers, President and Chief Executive Officer
- Rick Eiswirth, Chief Operating Officer and Chief Financial Officer
- Philip Ashman, Ph.D., Senior Vice President, European Managing Director
- Ken Green, Ph.D., Chief Scientific Officer, Global Head, Research & Development
- Dave Holland, Senior Vice President, Sales and Marketing

EU MANAGEMENT TEAM

- Philip Ashman, Ph.D., Senior Vice President, European Managing Director
- John Hall, MD, Senior Vice President, Medical Director Europe
- Philip Haldar, Vice President, European Marketing

LEAD PRODUCT

ILUVIEN is a long-term, sustained-release intravitreal implant. A single, tiny ILUVIEN implant is designed to deliver submicrogram levels of the corticosteroid, fluocinolone acetonide (FAC), for 36 months. Corticosteroids have a history of effective use in treating inflammation. ILUVIEN is injected into the back of the eye with an applicator that employs a 25-gauge needle, which allows for a self-sealing wound. In the two phase 3 clinical trials, known as the FAME™ Studies, the most frequently reported adverse drug reactions included cataract development and increased ocular pressure.

ILUVIEN IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

- ILUVIEN is contraindicated in patients with active or suspected ocular or periocular infections including most viral diseases of the cornea and conjunctiva including active epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, mycobacterial infections and fungal diseases.
- ILUVIEN is contraindicated in patients with glaucoma, who have cup to disc ratios of greater than 0.8.
- ILUVIEN is contraindicated in patients with known hypersensitivity to any components of this product.

WARNINGS AND PRECAUTIONS

- Intravitreal injections have been associated with endophthalmitis, eye inflammation, increased intraocular pressure, and retinal detachments. Patients should be monitored following the intravitreal injection.
- Use of corticosteroids may produce posterior subcapsular cataracts, increased intraocular pressure, glaucoma, and may enhance the establishment of secondary ocular infections due to bacteria, fungi, or viruses. Corticosteroids are not recommended to be used in patients with a history of ocular herpes simplex because of the potential for reactivation of the viral infection.
- Patients in whom the posterior capsule of the lens is absent or has a tear are at risk of implant migration into the anterior chamber.

ADVERSE REACTIONS

- In controlled studies, the most common adverse reactions reported were cataract development (ILUVIEN 82%; sham 50%) and intraocular pressure elevation of >10 mmHg (ILUVIEN 34%; sham 10%).
- Patients are advised to have follow-up eye examinations at appropriate intervals following treatment with ILUVIEN. For full prescribing information, log onto www.ILUVIEN.com.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see Full Prescribing Information below.

CORPORATE TIMELINE

- 2003: Alimera Sciences is founded by several former Novartis Ophthalmics executives.
- 2004: Soothe[®], an OTC product that provides relief to sufferers of dry eye, is launched.
- 2005: The FAME Study for ILUVIEN for the treatment of diabetic macular edema is initiated.
- 2006: Alimera gains FDA New Drug Application (NDA) approval for an Rx-to-OTC switch of an anti-allergy eye product called Alaway[®] ketotifen fumarate ophthalmic solution. Alaway is sold to Bausch and Lomb.
- 2007: Soothe is sold to Bausch and Lomb.
- 2010: Alimera Sciences goes public, trading under the NASDAQ symbol ALIM.
- 2012: ILUVIEN receives a positive opinion for approval of its Marketing Authorization Application (MAA) from seven European countries via the Decentralized Procedure (DCP).
- 2013: ILUVIEN is commercially launched in Germany and the United Kingdom.
- 2013: The United Kingdom's National Institute for Health and Care Excellence (NICE) published final guidance for ILUVIEN[®], clearing the path to patient availability through the National Health Service (NHS).
- 2014: Through the Repeat Use Process in Europe, Alimera receives a positive opinion to approve in 10 additional European countries and receives national marketing licenses in eight.
- 2014: ILUVIEN approved in the United States by the Food and Drug Administration (FDA).
- 2015: ILUVIEN commercially launched in the U.S. in late February.

FINANCIAL INFORMATION (as reported in the 10-Q dated November 14, 2014)

Alimera Sciences, Inc. is publicly traded on NASDAQ under the symbol ALIM. Net revenue for the three months ended September 30, 2014 increased by approximately \$1.6 million, or 211%, to approximately \$2.4 million compared to approximately \$760,000 for the three months ended September 30, 2013. The increase was attributable to increased adoption of ILUVIEN in Germany and the United Kingdom. As of September 30, 2014, Alimera had cash and cash equivalents of \$61.4 million, compared to \$12.6 million as of September 30, 2013.

LOCATIONS

Alimera Sciences, Inc.

6120 Windward Parkway, Suite 290
Alpharetta, GA 30005

T: +1 (678) 990 5740

F: +1 (678) 990 5744

Alimera Sciences Limited

Royal Pavilion
Wellesley Road
Aldershot Hampshire GU11 1PZ
United Kingdom

T: +44 (0)1252 354000

F: +44 (0) 1252 354012

CONTACTS

For press inquiries:

Audra Friis, Pascale Communications
917.519.9577
audra@pascalecommunications.com

For investor inquiries:

David Burke, The Ruth Group
646-536-7009
dburke@theruthgroup.com

For more information, visit www.alimerasciences.com.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ILUVIEN® safely and effectively. See full prescribing information for ILUVIEN.

ILUVIEN® (flucinolone acetonide intravitreal implant) 0.19 mg For Intravitreal Injection
Initial U.S. Approval: 1963

INDICATIONS AND USAGE

ILUVIEN contains a corticosteroid and is indicated for the treatment of diabetic macular edema in patients who have been previously treated with a course of corticosteroids and did not have a clinically significant rise in intraocular pressure. (1)

DOSAGE AND ADMINISTRATION

- For ophthalmic intravitreal injection. (2.1)
- The intravitreal injection procedure should be carried out under aseptic conditions. (2.2)
- Following the intravitreal injection, patients should be monitored for elevation in intraocular pressure and for endophthalmitis. (2.2)

DOSAGE FORMS AND STRENGTHS

Non-bioerodable intravitreal implant containing 0.19 mg flucinolone acetonide in a drug delivery system. (3)

CONTRAINDICATIONS

- Ocular or periocular infections (4.1)
- Glaucoma (4.2)
- Hypersensitivity (4.3)

WARNINGS AND PRECAUTIONS

- Intravitreal injections have been associated with endophthalmitis, eye inflammation, increased intraocular pressure, and retinal detachments. Patients should be monitored following the injection. (5.1)
- Use of corticosteroids may produce posterior subcapsular cataracts, increased intraocular pressure, glaucoma, and may enhance the establishment of secondary ocular infections due to bacteria, fungi, or viruses. (5.2)
- The implant may migrate into the anterior chamber if the posterior lens capsule is not intact. (5.3)

ADVERSE REACTIONS

In controlled studies, the most common adverse reactions reported were cataract development and increases in intraocular pressure. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Alimera Sciences, Inc. at 1-844-445-8843 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION

Revised: 12/2014

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information
2.2 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 4.1 Ocular or Periocular Infections
4.2 Glaucoma
4.3 Hypersensitivity

5 WARNINGS AND PRECAUTIONS

- 5.1 Intravitreal Injection-related Effects
5.2 Steroid-related Effects
5.3 Risk of Implant Migration

6 ADVERSE REACTIONS

- 6.1 Clinical Studies Experience
6.2 Postmarketing Experience

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ILUVIEN® (flucinolone acetonide intravitreal implant) 0.19 mg is indicated for the treatment of diabetic macular edema in patients who have been previously treated with a course of corticosteroids and did not have a clinically significant rise in intraocular pressure.

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information

For ophthalmic intravitreal injection.

2.2 Administration

The intravitreal injection procedure should be carried out under aseptic conditions, which include use of sterile gloves, a sterile drape, a sterile caliper, and a sterile eyelid speculum (or equivalent). Adequate anesthesia and a broad-spectrum microbicide should be given prior to the injection.

The injection procedure for **ILUVIEN** is as follows:

- The exterior of the tray should ***not*** be considered sterile. An assistant (non-sterile) should remove the tray from the carton and examine the tray and lid for damage. If damaged, do not use unit.

If acceptable, the assistant should peel the lid from the tray ***without touching the interior surface***.
- Visually check through the viewing window of the preloaded applicator to ensure that there is a drug implant inside.
- Remove the applicator from the tray with sterile gloved hands ***touching only the sterile interior tray surface and applicator***.

The protective cap on the needle should not be removed until the patient is ready to be injected.

Prior to injection, the applicator tip must be kept above the horizontal plane to ensure that the implant is properly positioned within the applicator.

- To reduce the amount of air administered with the implant, the administration procedure requires two steps. Before inserting the needle into the eye, push the applicator button down and slide it to the first stop (at the curved black marks alongside the button track). At the first stop, release the button and it should move to the UP position. If the button does not rise to the UP position, do not proceed with this unit.
- Optimal placement of the implant is inferior to the optic disc and posterior to the equator of the eye. Measure 4 millimeters inferotemporal from the limbus with the aid of calipers for point of entry into the sclera.
- Carefully remove the protective cap from the needle and inspect the tip to ensure it is not bent.
- Gently displace the conjunctiva so that after withdrawing the needle, the conjunctival and scleral needle entry sites will not align. Care should be taken to avoid contact between the needle and the lid margin or lashes. Insert the needle through the conjunctiva and sclera. To release the implant, while the button is in the UP position, advance the button by sliding it forward to the end of the button track and remove the needle. Note: Ensure that the button reaches the end of the track before removing the needle.
- Remove the lid speculum and perform indirect ophthalmoscopy to verify placement of the implant, adequate central retinal artery perfusion and absence of any other complications.

Following the injection, patients should be monitored for elevation in intraocular pressure and for endophthalmitis. Monitoring may consist of a check for perfusion of the optic nerve head immediately after the injection, tonometry within 30 minutes following the injection, and biomicroscopy between two and seven days following the injection. Patients should be instructed to report without delay any symptoms suggestive of endophthalmitis.

3 DOSAGE FORMS AND STRENGTHS

ILUVIEN is a non-bioerodable intravitreal implant in a drug delivery system containing 0.19 mg flucinolone acetonide, designed to release flucinolone acetonide at an initial rate of 0.25 µg/day and lasting 36 months.

4 CONTRAINDICATIONS

4.1 Ocular or Periocular Infections

ILUVIEN is contraindicated in patients with active or suspected ocular or periocular infections including most viral disease of the cornea and conjunctiva including active epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, mycobacterial infections and fungal diseases.

4.2 Glaucoma

ILUVIEN is contraindicated in patients with glaucoma, who have cup to disc ratios of greater than 0.8.

4.3 Hypersensitivity

ILUVIEN is contraindicated in patients with known hypersensitivity to any components of this product.

5 WARNINGS AND PRECAUTIONS

5.1 Intravitreal Injection-related Effects

Intravitreal injections, including those with **ILUVIEN**, have been associated with endophthalmitis, eye inflammation, increased intraocular pressure, and retinal detachments. Patients should be monitored following the intravitreal injection [see *Patient Counseling Information (17)*].

5.2 Steroid-related Effects

Use of corticosteroids including **ILUVIEN** may produce posterior subcapsular cataracts, increased intraocular pressure and glaucoma. Use of corticosteroids may enhance the establishment of secondary ocular infections due to bacteria, fungi, or viruses.

Corticosteroids are not recommended to be used in patients with a history of ocular herpes simplex because of the potential for reactivation of the viral infection.

5.3 Risk of Implant Migration

Patients in whom the posterior capsule of the lens is absent or has a tear are at risk of implant migration into the anterior chamber.

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse reactions associated with ophthalmic steroids including **ILUVIEN** include cataract formation and subsequent cataract surgery, elevated intraocular pressure, which may be associated with optic nerve damage, visual acuity and field defects, secondary ocular infection from pathogens including herpes simplex, and perforation of the globe where there is thinning of the cornea or sclera.

ILUVIEN was studied in two multicenter, randomized, sham-controlled, masked trials in which patients with diabetic macular edema were treated with either **ILUVIEN** (n=375) or sham (n=185).

Table 1 summarizes safety data available when the last subject completed the last 36 month follow up visit for the two primary **ILUVIEN** trials. In these trials, subjects were eligible for retreatment no earlier than 12 months after study entry. Over the three year follow up period, approximately 75% of the **ILUVIEN** treated subjects received only one **ILUVIEN** implant.

The most common ocular (study eye) and non-ocular adverse reactions are shown in Tables 1 and 2:

Table 1: Ocular Adverse Reactions Reported by ≥1% of Patients and Non-ocular Adverse Reactions Reported by ≥5% of Patients

Adverse Reactions	ILUVIEN (N=375) n (%)	Sham (N=185) n (%)
Ocular		
Cataract ¹	192/235 ² (82%)	61/121 ² (50%)
Myodesopsia	80 (21%)	17 (9%)
Eye pain	57 (15%)	25 (14%)
Conjunctival haemorrhage	50 (13%)	21 (11%)
Posterior capsule opacification	35 (9%)	6 (3%)
Eye irritation	30 (8%)	11 (6%)
Vitreous detachment	26 (7%)	12 (7%)
Conjunctivitis	14 (4%)	5 (3%)
Corneal oedema	13 (4%)	3 (2%)
Foreign body sensation in eyes	12 (3%)	4 (2%)
Eye pruritus	10 (3%)	3 (2%)
Ocular hyperaemia	10 (3%)	3 (2%)
Optic atrophy	9 (2%)	2 (1%)
Ocular discomfort	8 (2%)	1 (1%)
Photophobia	7 (2%)	2 (1%)
Retinal exudates	7 (2%)	0 (0%)
Anterior chamber cell	6 (2%)	1 (1%)
Eye discharge	6 (2%)	1 (1%)
Non-ocular		
Anemia	40 (11%)	10 (5%)
Headache	33 (9%)	11 (6%)
Renal Failure	32 (9%)	10 (5%)
Pneumonia	28 (7%)	8 (4%)

¹ Includes cataract, cataract nuclear, cataract subcapsular, cataract cortical and cataract diabetic in patients who were phakic at baseline. Among these patients, 80% of **ILUVIEN** subjects vs. 27% of sham-controlled subjects underwent cataract surgery.


² 235 of the 375 **ILUVIEN** subjects were phakic at baseline; 121 of 185 sham-controlled subjects were phakic at baseline.

Increased Intraocular Pressure

Table 2: Summary of Elevated IOP Related Adverse Reactions

Event	ILUVIEN (N=375) n (%)	Sham (N=185) n (%)
IOP elevation ≥ 10 mmHg from Baseline	127 (34%)	18 (10%)
IOP elevation ≥ 30 mmHg	75 (20%)	8 (4%)
Any IOP-lowering medication	144 (38%)	26 (14%)
Any surgical intervention for elevated intraocular pressure	18 (5%)	1 (1%)

Figure 1: Mean IOP during the study



Cataracts and Cataract Surgery

At baseline, 235 of the 375 **ILUVIEN** subjects were phakic; 121 of 185 sham-controlled subjects were phakic. The incidence of cataract development in patients who had a phakic study eye was higher in the **ILUVIEN** group (82%) compared with Sham (50%). The median time of cataract being reported as an adverse event was approximately 12 months in the **ILUVIEN** group and 19 months in the Sham group. Among these patients, 80% of **ILUVIEN** subjects vs. 27% of sham-controlled subjects underwent cataract surgery, generally within the first 18 months (Median Month 15 for both **ILUVIEN** group and for Sham) of the studies.

6.2 Postmarketing Experience

The following reactions have been identified during post-marketing use of **ILUVIEN** in clinical practice. Because they are reported voluntarily estimates of frequency cannot be made. The reactions, which have been chosen for inclusion due to either their seriousness, frequency of reporting, possible causal connection to **ILUVIEN**, or a combination of these factors, include reports of drug administration error and reports of the drug being ineffective.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

There are no adequate and well-controlled studies of **ILUVIEN** in pregnant women. Animal reproduction studies have not been conducted with flucinolone acetonide. Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. **ILUVIEN** should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.3 Nursing Mothers

Systemically administered corticosteroids are present in human milk and could suppress growth and interfere with endogenous corticosteroid production. The systemic concentration of flucinolone acetonide following intravitreal treatment with **ILUVIEN** is low [see Clinical Pharmacology (12.3)]. It is not known whether intravitreal treatment with **ILUVIEN** could result in sufficient systemic absorption to produce detectable quantities in human milk. Exercise caution when **ILUVIEN** is administered to a nursing woman.

8.4 Pediatric Use

Safety and effectiveness of **ILUVIEN** in pediatric patients have not been established.


8.5 Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

11 DESCRIPTION

ILUVIEN is a sterile non-bioerodable intravitreal implant containing 0.19 mg (190 mcg) flucinolone acetonide in a 36-month sustained-release drug delivery system. **ILUVIEN** is designed to release flucinolone acetonide at an initial rate of 0.25 µg/day. **ILUVIEN** is preloaded into a single-use applicator to facilitate injection of the implant directly into the vitreous. The drug substance is a synthetic corticosteroid, flucinolone acetonide.

The chemical name for flucinolone acetonide is (6α,11β, 16α)-6,9-difluoro-11,21-dihydroxy-16,17-[(1-methylethylidene)bis-(oxy)]-pregna-1,4-diene-3,20-dione. Its chemical structure is:



MW 452.50; molecular formula C₂₄H₃₀F₂O₆

Fluocinolone acetonide is a white or almost white, microcrystalline powder, practically insoluble in water, soluble in methanol, ethanol, chloroform and acetone, and sparingly soluble in ether.

Each **ILUVIEN** consists of a light brown 3.5mm x 0.37mm implant containing 0.19 mg of the active ingredient fluocinolone acetonide and the following inactive ingredients: polyimide tube, polyvinyl alcohol, silicone adhesive and water for injection.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Corticosteroids inhibit inflammatory responses to a variety of inciting agents. They inhibit edema, fibrin deposition, capillary dilation, leukocyte migration, capillary proliferation, fibroblast proliferation, deposition of collagen, and scar formation associated with inflammation.

Corticosteroids are thought to act by inhibition of phospholipase A₂ via induction of inhibitory proteins collectively called lipocortins. It is postulated that these proteins control biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting release of the common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A₂.

12.3 Pharmacokinetics

In a human pharmacokinetic study of **ILUVIEN**, fluocinolone acetonide concentrations in plasma were below the lower limit of quantitation of the assay (100 pg/mL) at all post-administration time points from Day 7 through Month 36 following intravitreal administration of a 0.2 mcg/day or 0.5 mcg/day fluocinolone acetonide insert.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been conducted to determine the carcinogenic potential or the effect on fertility of **ILUVIEN**.

Fluocinolone acetonide was not genotoxic *in vitro* in the Ames test (*S. typhimurium* and *E. coli*) and the mouse lymphoma TK assay, or *in vivo* in the mouse bone marrow micronucleus assay.

14 CLINICAL STUDIES

The efficacy of **ILUVIEN** was assessed in two three-year, randomized (2:1, active: sham), multicenter, double-masked, parallel-groups studies that enrolled patients with diabetic macular edema that had previously been treated with laser photocoagulation.

The primary efficacy endpoint in both trials was the proportion of subjects in whom vision had improved by 15 letters or more from baseline after 24 months of follow-up.

Table 3: Baseline BCVA (Letters)

	Study 1		Study 2	
	ILUVIEN (N=190)	Sham (N=95)	ILUVIEN (N=186)	Sham (N=90)
Mean (SD)	53 (13)	55 (11)	53 (12)	55 (11)
Median (Range)	57 (19-75)	58 (25-69)	56 (20-70)	58 (21-68)

Table 4: Visual Acuity outcomes at Month 24 (All randomized subjects with LOCF)

Study	Outcomes	ILUVIEN	Sham	Estimated Difference (95% CI)
1 ^a	Gain of ≥15 letters in BCVA (n (%))	51 (27%)	14 (15%)	12.1% (2.6%, 21.6%)
	Loss of ≥15 letters in BCVA (n (%))	26 (14%)	5 (5%)	8.4% (1.8%, 15.1%)
	Mean change from baseline in BCVA (SD)	3.7 (18.7)	3.2 (13.1)	1.8 (-2.8, 6.3)
2 ^b	Gain of ≥15 letters in BCVA (n (%))	57 (31%)	16 (18%)	13.0% (2.7%, 23.4%)
	Loss of ≥15 letters in BCVA (n (%))	22 (12%)	9 (10%)	1.8% (-5.9%, 9.6%)
	Mean change from baseline in BCVA (SD)	5.2 (18.0)	0.0 (15.6)	6.1 (1.4, 10.8)

^aStudy 1: **ILUVIEN**, N=190; Sham, N=95

^bStudy 2: **ILUVIEN**, N=186; Sham, N=90

Visual acuity outcomes by lens status (Phakic or Pseudophakic) at different visits are presented in Figure 2 and Figure 3. The occurrence of cataracts impacted visual acuity during the study. Patients who were pseudophakic at baseline achieved greater mean BCVA change from baseline at the Month 24 study visit.

Figure 2: Proportion of subjects with >=15 Letters Improvement from Baseline BCVA in the Study Eye





Figure 3: Mean BCVA Change from Baseline



The BCVA outcomes for the Pseudophakic and Phakic subgroups from Studies 1 and 2 at Month 24 are presented in Table 5.

Table 5: Visual Acuity outcomes at Month 24 (Subgroup for pooled data with LOCF)

Lens Status	Outcomes	ILUVIEN	Sham	Estimated Difference (95% CI)
^a Pseudophakic	Gain of ≥15 letters in BCVA (n (%))	39 (28%)	8 (13%)	15.4% (4.4%, 26.3%)
	Loss of ≥15 letters in BCVA (n (%))	7 (5%)	7 (11%)	-5.9% (-14.4%, 2.5%)
	Mean change from baseline in BCVA (SD)	7.1 (14.5)	1.5 (17.4)	5.6 (0.7, 10.6)
^b Phakic	Gain of ≥15 letters in BCVA (n (%))	69 (29%)	22 (18%)	11.1% (2.1%, 20.1%)
	Loss of ≥15 letters in BCVA (n (%))	41 (17%)	7 (6%)	11.6% (5.2%, 18%)
	Mean change from baseline in BCVA (SD)	2.8 (20.1)	1.8 (12.6)	1 (-2.5, 4.4)

^aPseudophakic: **ILUVIEN**, N=140; Sham, N=64

^bPhakic: **ILUVIEN**, N=236; Sham, N=121

16 HOW SUPPLIED/STORAGE AND HANDLING

ILUVIEN[®] (fluocinolone acetonide intravitreal implant) 0.19 mg is supplied in a sterile single use preloaded applicator with a 25-gauge needle, packaged in a tray sealed with a lid inside a carton.

NDC 68611-190-02

Storage: Store at 15°-30° C (59°-86° F).

17 PATIENT COUNSELING INFORMATION

Steroid-related Effects

Advise patients that a cataract may occur after treatment with **ILUVIEN**. If this occurs, advise patients that their vision will decrease, and they will need an operation to remove the cataract and restore their vision.

Advise patients that they may develop increased intraocular pressure with **ILUVIEN** treatment, and the increased IOP may need to be managed with eye drops, or surgery.

Intravitreal Injection-related Effects

Advise patients that in the days following intravitreal injection of **ILUVIEN**, patients are at risk for potential complications including in particular, but not limited to, the development of endophthalmitis or elevated intraocular pressure.

When to Seek Physician Advice

Advise patients that if the eye becomes red, sensitive to light, painful, or develops a change in vision, they should seek immediate care from an ophthalmologist.

Driving and Using Machines

Inform patients that they may experience temporary visual blurring after receiving an intravitreal injection. Advise patients not to drive or use machines until this has been resolved.

Manufactured for:

Alimera Sciences, Inc.
6120 Windward Parkway
Alpharetta, GA 30005

Patented. See: www.alimerasciences.com

ALIMERA
SCIENCES