Rivaroxaban in Venous and Arterial Thromboembolism (VAT)

What is Venous and Arterial Thromboembolism (VAT)?

Venous and arterial thromboembolism (VAT) is caused when all, or part of a blood clot breaks away (an embolus) and is moved by the blood stream to block a vein or artery. This obstruction can result in damage to vital organs, because the tissue beyond the blockage no longer receives nutrients and oxygen.

VAT is responsible for increasing morbidity and mortality across a broad range of acute and chronic blood-clotting conditions.

In the EU, more people die from blood clots than from AIDS, breast cancer, prostate cancer and traffic accidents combined¹

Venous Thromboembolism (VTE)

VTE is the third most common cardiovascular disease worldwide² and is the most common avoidable cause of hospital death³

VTE encompasses two conditions: Deep vein thrombosis (DVT) is a blood clot that forms in the veins that lie deep within the muscles, usually in the leg or pelvis. If all or part of the DVT breaks off and the blood clot moves to block a vessel in the lungs, it is known as a **pulmonary embolism (PE)**⁴, which can be rapidly fatal.

Symptoms of DVT include: Pain, swelling, redness of the affected area, usually the leg, and the dilation of surface veins; the skin may also be warm to the touch Symptoms of PE include: Acute shortness of breath, chest pain, rapid heart rate and light headedness; some people may also cough blood

Who is at Risk of VTE?

- Patients undergoing surgery such as hip or knee replacement or major orthopaedic surgery or major surgery for cancer
- Patient-related risk factors include inherited thrombophilia, advanced age, obesity, prior VTE and varicose veins⁴
- Patients admitted to hospital for an acute medical condition

Arterial Thromboembolism

 Arterial thromboembolism occurs when oxygenated blood flow from the heart to another part of the body, via an artery, is interrupted by a blood clot

Pulmonarv

Embolism (PE)

Venous Thromboembolism (VTE)

- If this occurs in an artery supplying blood to the brain it can lead to a stroke, which can be severely debilitating or fatal
- If it occurs in a coronary artery, it can lead to acute coronary syndrome (ACS), which includes conditions such as myocardial infarction (heart attack) and unstable angina

Who is at Risk for Arterial Thromboembolism?

- Patients with atrial fibrillation (AF) and those who have survived a previous episode of ACS
- The essential underlying condition for ACS is the build-up of plaque in the inner walls of coronary arteries that narrows the arteries. This process is called atherosclerosis. When atherosclerotic plaque ruptures or erodes it could lead to a sudden and critical reduction of blood flow

How is VAT Prevented and Treated?

VAT Encompasses Two Serious Conditions

Deep Vein

Thrombosis (DVT)

VAT requires active or preventative treatment to avoid potentially serious or fatal patient outcomes.

Stroke

Arterial Thromboembolism

Heart


Attack



However, these older anticoagulants are associated with significant drawbacks for the patient such as regular blood coagulation monitoring or the need for injections.

Novel oral anticoagulants (OACs) can overcome the limitations of older anticoagulants to prevent and/or treat these potentially deadly blood clots.

Benefits of novel OACs include⁵:



Low risk of drug-drug interactions



Rivaroxaban is an effective, novel oral anticoagulant developed to prevent and treat dangerous blood clots by selectively targeting Factor Xa, an enzyme which acts at the pivotal stage in bloodclotting. With more than 10 million patients treated* over six years across five indications in seven areas of use, rivaroxaban is the most prescribed novel OAC in the world^{5,6}

IMS Health MIDAS, Database. Monthly Sales July 2014



Rivaroxaban in Venous and Arterial Thromboembolism (VAT) - Continued



About Rivaroxaban

Rivaroxaban is the most broadly indicated and most prescribed novel OAC⁶ and is marketed under the brand name Xarelto[®]. Rivaroxaban is approved for five indications across seven distinct areas of use, protecting patients across more venous and arterial thromboembolic (VAT) conditions than any other novel OAC:



The prevention of stroke and systemic embolism in adult patients with nonvalvular atrial fibrillation (AF) with one or more risk factors




The prevention of venous thromboembolism (VTE) in adult patients undergoing elective hip replacement surgery



replacement surgery

The prevention of VTE in adult patients undergoing elective knee



The treatment of pulmonary embolism (PE) in adults*



The prevention of recurrent DVT and PE in adults



The prevention of atherothrombotic events (cardiovascular death, heart attack or stroke) after an acute coronary syndrome in adult patients with elevated cardiac biomarkers when co-administered with acetylsalicylic acid (ASA) alone or with ASA plus clopidogrel or ticlopidine

Whilst licences may differ from country, across all indications rivaroxaban is approved in more than 125 countries. Rivaroxaban was discovered by Bayer HealthCare, and is being jointly developed with Janssen Research & Development, LLC. Rivaroxaban is marketed outside the U.S. by Baver HealthCare and in the U.S. by Janssen Pharmaceuticals, Inc. (a Johnson & Johnson Company).

Anticoagulant medicines are potent therapies used to prevent or treat serious illnesses and potentially life threatening conditions. Before initiating therapy with anticoagulant medicines, physicians should carefully assess the benefit and risk for the individual patient. Responsible use of rivaroxaban is a very high priority for Bayer, and the company has developed a Prescribers Guide for physicians and a 'Xarelto' Patient Card for patients to support best practice.

To learn more, please visit https://prescribe.xarelto.com To learn more about thrombosis, please visit www.thrombosisadviser.com

To learn more about VAT, please visit www.VATspace.com To learn more about 'Xarelto', please visit www.xarelto.com

References

1) Cohen AT, Agnelli G, Anderson FA, et al. Thromb Haemost. 2007;98.(4)756-764 2) Naess IA, Christiansen SC, Romundstad P, et al. J Thromb Haemost. 2007;5(4):692-699 3) Geerts WH, Bergqvist D, Pineo GF, et al. Chest. 2008;133,(6 Suppl)381S-453S 4) Patient UK. Deep vein thrombosis. Available at: http://www.patient.co.uk/health/Deep-Vein-Thrombosis.htm Accessed January 2015 5) Xarelto Summary of Product Characteristics as approved by the European Commission 6) IMS Health MIDAS, Database. Monthly Sales July 2014 *Rivaroxaban is not recommended as an alternative to unfractionated heparin in patients with PE who present hemodynamic instability or who may receive thrombolysis or pulmonary embolectomy